

Family planning in the slums of Nairobi, Kenya

Planning familial dans les bidonvilles de Nairobi, Kenya

[Donatien Beguy, Alex C. Ezeh, Blessing U. Mberu, Jacques B.O. Emina](#)

In Kenya, rapid population growth has occurred amidst poor urban governance and limited employment opportunities, leading to widespread urban poverty that is concentrated in informal settlements or slums. In Nairobi, the capital city, it is estimated that between 60 and 70% of the residents live in slums or slum-like conditions (APHRC 2014). Slum-dwellers have limited access to health and reproductive health services and they have comparatively poor sexual and reproductive health outcomes (Zulu et al. 2008). Within urban areas in Kenya, unmet need for contraception is greater among the poorest women, and their childbearing is often mistimed or unwanted (Ezeh et al. 2010). However, it is suggested that the gap between poor and rich in the use of modern contraception in urban Kenya is narrowing, and that family planning (FP) services may be increasingly reaching the urban poor (Fotso et al. 2013).

Using data collected from representative samples of the slum population in Nairobi in 2000 and 2012, we recently examined changes in the use of modern contraception in Nairobi slums over the period (Beguy et al 2017).

Our findings

Overall, there was a dramatic 19-percentage-point increase in the use of modern contraception among married women between 2000 (35%) and 2012 (54%), with a substantial improvement among hitherto disadvantaged groups, i.e. poor urban women. This is consistent with the substantial increase in modern contraceptive prevalence rate (CPR) over the last decade in Kenya, from 32% in 2003 to 53% in 2014, with the public sector being the major provider of contraceptive methods. While in 2000 there were significant differentials in the CPR by various socio-demographic and socioeconomic characteristics, these associations had nearly disappeared in 2012. For example, in 2000, women with secondary or higher education were much more likely to be using modern methods than those with no education (43% vs. 13%). By 2012, usage had increased in both groups, and the differences had considerably shrunk (54% and 33%, respectively; see Table 1).

TABLE 1. Percent of women using modern contraception by selected characteristics in the slums of Nairobi (Kenya) in 2000 and 2012

Characteristic	2000		2012		Absolute change	Relative change
	Percent	N	Percent	N		
Age group						
15-19	28.5	254	45.8	356	18.3	64.2
20-24	42.2	340	58.2	467	17.0	40.2
25-29	26.7	161	45.8	277	20.1	67.7
Ethnic group						
Kalenjin	24.0	133	53.0	270	29.0	117.0
Kisumu	32.7	167	47.3	250	14.6	45.0
Luhya	33.0	209	51.5	307	18.5	56.0
Luo	31.1	67	49.2	183	18.1	58.2
Other	33.5	70	58.5	273	25.0	86.1
Women's education level						
No education	12.8	74	32.8	74	19.8	162.7
Primary	31.0	364	53.8	655	22.8	73.6
Secondary+	44.3	287	62.8	389	18.5	41.7
Religion						
Catholic	31.5	165	55.5	305	24.0	69.7
Protestant/Other Christian	28.7	468	52.3	470	23.6	68.2
Muslim	24.4	23	42.1	53	17.7	72.8
No religion	20.0	2	45.1	8	25.1	250.0
Other religions	33.2	7	43.8	8	10.6	31.8
Health status						
Poor	25.8	150	48.8	388	23.1	66.6
Moderate	32.4	239	50.7	488	18.3	56.1
Good	41.8	237	58.2	478	16.4	39.0
Women's employment status						
Unemployed	22.4	220	30.8	342	8.4	37.5
Employed	43.8	339	66.8	672	23.1	52.7
Number of living children						
0	6.1	78	17.2	87	11.2	182.8
1	36.0	342	52.8	364	17.8	49.4
2	44.2	370	52.5	366	8.3	18.8
3	11.1	122	16.2	121	5.1	45.9
4	4.3	72	8.2	112	3.9	90.7
5+	1.0	60	3.8	78	2.8	280.0
Women's social inequality experience						
None	17.2	127	54.1	1,155	36.9	45.4
One or more	23.7	88	47.4	157	23.7	100.2
Had about FP at a health facility						
Did not visit	26.0	129	48.8	381	22.8	79.2
Visited and used about FP	52.3	348	62.7	467	10.4	20.1
Visited but did not use about FP	21.9	292	34.5	473	12.6	58.0
Family preferences						
Wanted additional child	30.4	1135	48.2	586	17.8	57.4
Wanted no more	38.4	749	67.8	828	29.4	68.2
Undecided	48.9	68	31.7	68	-17.2	-35.0
Total	34.3	861	52.8	1,244	18.5	54.1

SOURCE: Authors' calculations on data from the Nairobi Urban Reproductive Health Survey (URHS) 2000 and 2012.

Overall, data suggests that the observed increase in modern CPR among women of reproductive age in Nairobi slums between 2000 and 2012 was overwhelmingly due to a change in their reproductive behavior with respect to use of modern contraception. This is consistent with findings from a recent study which demonstrated that changes in behavior explained much of the increase in modern CPR in 27 countries in sub-Saharan Africa, including Kenya (Emina et al. 2014).

Family planning programs apparently work

The change observed in Nairobi's slums was probably due to the revival of family planning (or FP) programs in Kenya over the past few years. To address the significant disparities in contraceptive use and unmet need for FP in the country, the government of Kenya sought to eliminate stockouts of contraceptive commodities and remove financial barriers to family planning services by introducing, in 2006, a budget line for contraceptive commodities in the Ministry of Health's annual budget. This has significantly narrowed most of the gaps in CPR since 2008-09. Other programs were set up to expand the reach and quality of FP programs among disadvantaged groups, including women from urban slums. For example, the Gates Foundation-funded Kenya Urban Reproductive Health Initiative (Tupange) was implemented between 2009 and 2015 to raise the CPR by at least 20 percentage points in poor urban settlements in five cities: Nairobi, Kisumu, Mombasa, Kakamega, and Machako.

Our analysis shows that more effective meeting of fertility preferences, particularly among women who want to cease childbearing, has contributed to much of the increase in use of modern contraception among women in Nairobi slums. Those who want no more children have the highest levels of contraceptive use: over 60% in 2012, reflecting a 56% increase over the period 2000-2012.

In addition, compositional change also plays a role in increasing CPR in Nairobi slums. For instance, the increase from 34% to 46% in the proportion of women with at least secondary education and the fact that these women also have the

highest levels of contraceptive use explains much of the change in modern contraceptive use between 2000 and 2012. This finding corroborates previous evidence showing women's education to be a key determinant of modern contraceptive use, and suggests that, even in slum settlements, education may have a powerful effect in bringing about social change.

Data also show that as child survival has improved over time, the demand for contraception to space or limit childbearing has increased among women in Nairobi slums.

Although we were unable to assess changes in the quality of reproductive health services, the proportion of women reporting that they were informed about FP when they visited a health facility increased considerably between 2000 and 2012. Our analysis also shows significant behavioral effects, suggesting that programs may be adopting strategies to ensure that women attending health facilities are provided with appropriate information to assist those who wish to adopt FP methods. For example, Tupange activities include training health care providers, FP orientation for all facility staff, and integration of FP services with other maternal and child health and HIV/AIDS services (Speizer et al. 2013). Also, the program's social franchising of private providers has helped expand provision of FP services, in conjunction with its advocacy for increased resources for FP to improve commodity security and create a conducive policy environment. Other activities include working with community health workers to promote and distribute FP methods; community awareness and educational activities with local leaders, women, and youth.

Conclusion

This research suggests that in Nairobi slums, the largest gains in FP uptake between 2000 and 2012 were made among previously disadvantaged groups. Much higher increases among previously deprived groups have reduced inequity in access to contraceptive methods, constituting the main force behind the increase in overall contraceptive use. This may be the result of efforts by the government of Kenya and its development partners to reduce the disparities in uptake and use of FP services by targeting the poorest segments of the Kenyan population, such as those living in urban slums. This success is also an indication that such programs need to be sustained and scaled-up across the country.

References

APHRC – African Population and Health Research Center (2014). *Population and Health Dynamics in Nairobi's Informal Settlements: Report of the Nairobi Cross-sectional Slums Survey (NCSS) 2012*. Nairobi: African Population and Health Research Center.

Beguy D., Ezeh A.C., Mberu B.U., Emina J.B.O. (2017) Changes in Use of Family Planning among the Urban Poor: Evidence from Nairobi Slums. *Population and Development Review*, supplement chapter. DOI: 10.1111/padr.12038.

Ezeh AC, Kodzi I, Emina J (2010). Reaching the Urban Poor with Family Planning Services. *Studies in Family Planning*. 41(2):109-116.

Fotso J, Speizer I, Mukiira C, Kizito P, Lumumba V. 2013. Closing the poor-rich gap in contraceptive use in urban Kenya: are family planning programs increasingly reaching the urban poor? *International Journal for Equity in Health*. 12(1):71.

Emina JBO, Chirwa T, Kandala N-B (2014). Trend in the use of modern contraception in sub-Saharan Africa: does women's education matter? *Contraception*.90(2):154-161.

Speizer IS, Corroon M, Calhoun L, et al. (2014). Demand generation activities and modern contraceptive use in urban areas of four countries: a longitudinal evaluation. *Global health, science and practice*. 2(4):410-426.

Zulu EM, Dadoo FN-A, Ezeh AC (2002). Sexual risk-taking in the slums of Nairobi, Kenya, 1993-98. *Population Studies*. 56(3):311-323.