

Towards a better identification of older people at risk of depression

Omar Paccagnella | September 9, 2021



Depression is one of the most common mental disorders in later life. Omar Paccagnella suggests a new approach to classify older individuals into distinct subgroups, based on different combinations of depressive and emotional states. This allows scholars to go beyond the traditional depression / non-depression dichotomy.

Introduction

Depression is one of the most common mental disorders affecting older people. It often coexists with other medical conditions and physical impairments (Blazer, 2003), and affects women more than men.

Later-life depression is characterized by a broad range of symptoms. The identification of substantial and meaningful subgroups of depressive symptom profiles has important implications for research, public health policies and clinical practice, offering scope for developing customized intervention and treatment programmes.

A range of screening methods are available for detecting depressive disorders and can be divided into two broad categories: tools to collect information on diagnostic criteria that best reflect diagnoses of mental disorders; and tools to measure more generic factors (psychological distress) and the presence of certain symptoms or emotional disorders, with scores and cut-off points (usually just one) to classify individuals with or without depression or mental health disorders.

Going beyond dichotomisation

A more nuanced classification based on the number and type of reported items in a depression scale would be preferable to a simple dichotomisation of 'depressed' versus 'non-depressed'.

To this end, we tried an alternative approach based on two statistical techniques:

1. a *latent class analysis* to estimate the individual probability of belonging to a specific category (with respect to risk of depression) and then
2. a *factor analysis*, to reduce the number of clusters (Gennaro et al., 2021).

We analysed data collected in the sixth wave of the Survey of Health, Ageing and Retirement in Europe (SHARE), carried out in 2015 (Börsch-Supan, 2017).

We focused on the 12 items of the EURO-D scale of depressive symptoms, developed by the EURODEP Concerted Action Programme: depressed mood, pessimism, wishing death, guilt, sleep, interest, irritability, appetite, fatigue, concentration, enjoyment, tearfulness (Table A1). EURO-D is a symptom-oriented scale, with scores ranging between 0 and 12. Scores above 3 are customarily assumed to reveal depression (Prince et al., 1999).

Table A1: List of questions and response options of the EURO-D scale in SHARE

Item	Main question	Response options
Depressed mood	In the last month, have you been sad or depressed?	1. Yes 5. No
Pessimism	What are your hopes for the future?	1. Any hopes mentioned 2. No hopes mentioned
Suicidality	In the last month, have you felt that you would rather be dead?	1. Any mention of suicidal feelings or wishing to be dead 2. No such feelings
Guilt	Do you tend to blame <u>yourself</u> or feel guilty about anything?	1. Obvious excessive guilt or self-blame 2. No such feelings 3. Mentions guilt or self-blame, but it is unclear if these constitute obvious or excessive guilt or self-blame ^(a)
Sleep	Have you had trouble sleeping recently?	1. Trouble with sleep or recent change in pattern 2. No trouble sleeping
Interest	In the last month, what is your interest in things?	1. Less interest than usual mentioned 2. No mention of loss of interest 3. Non-specific or uncodeable response ^(b)
Irritability	Have you been irritable recently?	1. Yes 5. No
Appetite	What has your appetite been like?	1. Diminution in desire for food 2. No diminution in desire for food 3. Non-specific or uncodeable response ^(c)
Fatigue	In the last month, have you had too little energy to do the things you wanted to do?	1. Yes 5. No
Concentration	How is your concentration? For example, can you concentrate on a television programme, film or radio programme? ^(d)	1. Difficulty in concentrating on entertainment 2. No such difficulty mentioned
Enjoyment	What have you enjoyed doing recently?	1. Fails to mention any enjoyable activity 2. Mentions ANY enjoyment from activity
Tearfulness	In the last month, have you cried at all?	1. Yes 5. No

Additional questions

(a) So, for what do you blame yourself? [Possible answers: Example(s) given do or do not constitute obvious excessive guilt or self-blame]

(b) So, do you keep up your interests? [Possible answers: yes/no]

(c) So, have you been eating more or less than usual? [Possible answers: less/more/neither]

(d) Can you concentrate on something you read? [Possible answers: with/without difficulty].

In our sample (64,716 individuals aged 50 years and over, living in 18 countries: Austria, Belgium, Croatia, Czech Republic, Denmark, Estonia, France, Germany, Greece, Italy, Israel, Luxembourg, Poland, Portugal, Slovenia, Spain, Sweden and Switzerland) about 27% of respondents (20% of men, and 33% of women) are depressed according to the EURO-D scale (Table 2).


Table 2. Classification and total number of reported depressive symptoms according to the EURO-D scale, by gender (European respondents aged 50 years and over, 2015)

EURO-D classification	Number of reported EURO-D items	Proportion (%)		
		Men	Women	All
NON DEPRESSED	0	28.0	17.6	22.1
	1	23.5	18.5	20.7
	2	17.2	16.5	16.7
	3	11.8	14.1	13.1
	Total	80.5	66.6	72.6
DEPRESSED	4	7.7	11.1	9.6
	5	4.9	8.0	6.7
	6	3.0	5.7	4.5
	7	1.8	3.6	2.9
	8	1.1	2.4	1.8
	9	0.6	1.5	1.1
	10	0.3	0.7	0.5
	11	0.1	0.3	0.2
12	<0.1	0.1	0.1	
Total	19.5	33.4	27.4	

Source: SHARE data. See Gennaro et al. (2021).

There is substantial heterogeneity in the number of reported symptoms: some are frequent, such as depressed mood, sleep impairment, irritability, fatigue and (only for women) tearfulness, while some are rare, such as suicidality and guilt (Figure 1). Among women classified as non-depressed by EURO-D, only one-fourth reported no symptoms at all, a proportion that rises to about one-third among men.

Figure 1. Distribution of reported depression symptoms, by gender and EURO-D classification (European respondents age 50 years and over, 2015)



Source: SHARE data. See Gennaro et al. (2021).

Main results

Our more refined approach leads to the identification of seven (main) categories of depressive and emotional states, from very low risk of depression to extreme depression (Table 3). These groups are characterised by different combinations of symptoms. As the attribution of individuals to one of the seven clusters is guided by probabilistic criteria, our solution does not necessarily group individuals with the same number of symptoms: rather, it identifies common patterns in these symptoms.

Table 3. Depression categories according to our approach (European respondents age 50 years and over, 2015)


Category	Name	Size (%)		
		Men	Women	All
1	Very low risk of depression	48.3	34.4	40.4
2	Low risk of depression	3.2	1.9	2.5
3	Moderate risk of depression	13.8	21.0	17.9
4	High risk of depression	29.0	34.2	32.0
5	Depressed	2.9	3.2	3.0
6	Severely Depressed	2.3	4.2	3.4
7	Extremely depressed	0.5	1.1	0.8

Source: SHARE data. See Gennaro et al. (2021).

About 40% of the people in our sample are at a very low risk of depression: they present a similar combination of symptoms, in particular between zero and two reported items at most. The last three categories are characterised by individuals with a high probability of reporting several depressive and emotional problems. One-third of respondents fall into the high-risk-of-depression category, i.e. they are not depressed, strictly speaking, but might well become so in the future. Ex-post analyses of the composition of this group reveal that unhealthy respondents are overrepresented (for instance, almost 60% of them report two or more chronic diseases), as are women (more than 60%).

A large proportion of individuals classified as depressed under the EURO-D system falls into one of the clusters with some risk of depression (Figure 2). Under our approach, however, a non-trivial proportion of older adults classified as non-depressed by EURO-D are identified as being at some risk of depression. Almost all respondents in the three most severe depression categories are also classified as depressed by the EURO-D scale. Gender differences are particularly evident.

Figure 2. Distribution of the categories of our approach within each group of respondents classified as depressed or not according to the EURO-D scale, by gender (European respondents age 50 years and over, 2015)



Source: SHARE data. See Gennaro et al. (2021).

As a first step, we applied our approach to cross-sectional data only. However, given the dynamic nature of depression, it would be very interesting to follow developmental trajectories or patterns of change over time. This should be possible with our approach, using Latent Class Growth Models as a first step. Hopefully, then, more on this in the near future.

References

- Blazer DG (2003). Depression in late life: review and commentary. *Journals of Gerontology* 58A, 249-265
- Börsch-Supan A (2017). Survey of Health, Ageing and Retirement in Europe (SHARE) Wave 6. Release version 6.0.0. SHARE-ERIC (Data set). doi:10.6103/SHARE.w6.600
- Gennaro C, Paccagnella O, Zaninotto P (2021). A model-driven approach to better identify older people at risk of depression. *Ageing & Society* 41, 339-361
- Prince M, Reischies F, Beekman AT, Fuhrer R et al. (1999). Development of the EURO-D scale - a European Union initiative to compare symptoms of depression in 14 European centres. *British Journal of Psychiatry* 174, 330-338